

# **Regional Stratification and Shear of the Various Streams Feeding the Philippine Straits – ESR Component**

Amy Ffield

Earth & Space Research, 290 Clausland Mountain Road, Upper Grandview, NY 10960-4113  
phone: (845) 353-1733 fax: (845) 353-1733 email: [ffield@esr.org](mailto:ffield@esr.org)

Award Number: N0001406C0578  
<http://www.esr.org>

## **LONG-TERM GOALS**

To determine the impact of the regional oceanographic and atmospheric mean and variable conditions on the Philippine Archipelago strait dynamics.

## **OBJECTIVES**

To quantify the monsoonal and interannual oceanographic stratification and shear boundary conditions in the Philippine Archipelago region.

## **APPROACH**

Regional CTD and Lowered ADCP [CTD/LADCP] study in cooperation with Arnold Gordon (Lamont-Doherty Earth Observatory).


## **WORK COMPLETED**

Background data analysis on the Philippine Archipelago region has been completed in preparation for the field experiment. (At the time of writing this report the P.I. has not yet received funds for the project.)


## **RESULTS**

*Sea Surface Temperature:* Analysis of the OI Infrared Satellite SST reveals that the SST in the Philippine Archipelago region has significant variability (Figure 1). This variability may be mirrored in the deeper ocean possibly impacting strait dynamics. For example, in 1987 the annual range in SST is quite large:  $\sim 4.5^{\circ}\text{C}$ , but in 1988 the annual SST range is only  $\sim 1.5^{\circ}\text{C}$  (**Figure 2**); these SST differences may reflect quite different sub-surface ocean stratifications that could impact strait dynamics.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2006</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2006 to 00-00-2006</b>	
4. TITLE AND SUBTITLE <b>Regional Stratification and Shear of the Various Streams Feeding the Philippine Straits - ESR Component</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Earth &amp; Space Research, 290 Clausland Mountain Road, Upper Grandview, NY, 10960-4113</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>5</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			




**Figure 1.** The average February SST (a) and the interannual SST contrasts between the warmest and coolest Februaries (b) in the Philippine region.




**Figure 2.** SST time series for the Mindoro Strait / northern Sulu Sea region in the Philippine Archipelago.


*Sea Level/Tide Gauges:* Multiple sea level/tide gauges are in operation in the Philippine Archipelago region (**Figure 3**). The gauges capture the tidal variability (**Figure 4**), as well as significant mean sea level height differences across the basins (**Figure 5**) that can be used to characterize the background conditions impacting strait dynamics.



*Figure 3. Map of the sea level height station locations in the Philippine Archipelago region. Data may not be available from the gauges at Jolo and Puerto Princesa at the time of the field experiment.*




*Figure 4. Sea level height time series for the gauges in the Philippine Archipelago region.*




**Figure 5.** *Sea level height difference between the Manila and Sandakan tide gauges, potentially representing the variability in the cross-channel sea level height difference of Mindoro Strait.*

*Barotropic Tides:* The altimeter-derived barotropic modeled tide heights on either side of the Mindoro Strait (**Figure 6**) reveal spring tide ranges as large as 2 m (**Figure 7**). The difference in the barotropic tides across the straits may also be important to strait dynamics; in this case, the predicted barotropic tide height difference across the Mindoro Strait varies from as low as 0.02 m during neap tides up to 0.25 m during spring tides (**Figure 8**).




**Figure 6.** *Map of the predicted barotropic tide height locations on either side of Mindoro Strait.*





*Figure 7. Time series of the predicted barotropic tide heights on either side of Mindoro Strait.*

*Figure 8. Time series of the difference in predicted barotropic tide height across Mindoro Strait.*



## IMPACT/APPLICATIONS

The preparatory analysis will aid in the final design of the field experiment.